

Introduction

Let us consider the initial-boundary problem for the Davey-Stewartson-II system of equations:

$$iu_t + u_{xx} - u_{yy} = \gamma|u|^2u + \lambda u\varphi_x, \quad \varphi_{xx} + \varphi_{yy} = \mu(|u|^2)_x, \quad (1)$$

$$u(x, y, t)|_{t=0} = u_0(x, y); \quad \nabla\varphi \rightarrow 0, \quad x^2 + y^2 \rightarrow \infty. \quad (2)$$

The Davey-Stewartson-II system of equations is the two-dimensional generalization of the usual nonlinear Schrodinger equation. Systems of the form (1) were first derived by Davey and Stewartson in the shallow-water limit for the surface waves in fluid, [1]. Physical applications also include plasma physics and nonlinear optics. In the above $u(x, y, t)$ is the amplitude of a surface wave packet and $\varphi(x, y, t)$ is the velocity potential of mean flow interacting with the surface wave. It is worth mentioning that some of Davey-Stewartson systems of equations are integrable, but many of them are not integrable. The existence and uniqueness results with respect to the initial data (in L_2, H^1, H^2) were recently obtained in the sufficiently wide classes of generalized solutions, [2]. Our notice is centred on existence of classical solutions of the initial-boundary problem for the more generally Davey-Stewartson-II system of equations.

The existence and uniqueness theorems

Fourier transform with respect to spatial variables lies in the base of proof of solvability of the initial-boundary problem (1)-(2). By using this traditional approach to the proof of existence theorems, we may pass from the system of partial differential equations (1) to a single integral equation in terms of the Fourier transform $\hat{u}(k, m, t)$ of $u(x, y, t)$:

$$\hat{u}(k, m, t) = \hat{u}_0(k, m)e^{i(m^2 - k^2)t} - i \int_0^t e^{i(k^2 - m^2)(\tau - t)} L(\hat{u}(k, m, \tau)) e^{2i(k^2 - m^2)\tau} d\tau. \quad (3)$$

where the operator L is defined by the following formula:

$$L(\hat{u}) = \gamma \hat{u} \star \hat{u}^* \star \hat{u} + \lambda \mu \hat{u} \star \left[\frac{k^2}{k^2 + m^2} (\hat{u} \star \hat{u}^*) \right].$$

Here the sign $\star$ is a convolution with respect to (k, m) .

Theorem 1. Let $u_0(x, y) \in H^p$ provided $p > 1$. There exists a positive $T > 0$ such that $\forall t \in [0, T]$ the initial-boundary problem (1)-(2) has a unique solution such that: $u, \nabla\varphi \in C^1([0, T]; H^p)$.

The basic idea of the proof is now to use a contraction mapping method in a suitable scale of Banach spaces. In these Banach spaces integral operator from (3) should be bounded and Lipschitzian. By analogy with [3,4], we introduce the scale of Banach spaces $C([0, T]; H^p)$ of Fourier transforms $\hat{u}(k, m, t)$, which are continuous with respect to (k, m) , $t \in [0, T]$ and decreasing at the infinity, with the following norms:

$$\|\hat{u}(k, m, t)\| = \sup_{t \in [0, T]} \|(1 + |k| + |m|)^p \hat{u}(k, m, t)\|_{L_2}.$$

In view of nonlinearity of L operator, our Banach space must be an algebra with respect to the convolution operation. This problem was solved provided $p > 1$. It turns out that in this case the convolution is a Lipschitzian operator in Sobolev space H^p . Then if we choose a suitable T such that the integral operator from (3) would be contraction, the solvability of the equation (3), and so of the problem (1)-(2) follows from the Ovsjannikov's results for the equations in Banach spaces, [5].

This result is directly applicable to the more general systems of equations in multidimensions. Namely, we consider the following initial-boundary problem for the system of partial differential equations:

$$i\partial_t u + \omega(-i\partial_{x_1}, \dots, -i\partial_{x_n})u = g(u, u^*) + \bar{h}(u, u^*)\nabla\varphi, \quad P\varphi = \nabla\bar{f}(u, u^*), \quad (4)$$

$$u(x_1, \dots, x_n, t)|_{t=0} = u_0(x_1, \dots, x_n); \quad \nabla\varphi \rightarrow 0, \quad x_1^2 + \dots + x_n^2 \rightarrow \infty, \quad (5)$$

where P is a second-order differential operator with the constant coefficients and $\omega(k_1, \dots, k_n)$ is a polinom with respect to $k_1, \dots, k_n$.

Theorem 2. Let 1) functions $\bar{f}, g, \bar{h}$ are analytical with respect to all arguments such that: $\bar{f}(0, 0) = g(0, 0) = \bar{h}(0, 0) = 0$; 2) initial function $u_0(x_1, \dots, x_n) \in H^p$ provided $p > n/2$. There exists a positive $T > 0$ such that $\forall t \in [0, T]$ the initial-boundary problem (4)-(5) has a unique solution such that: $u, \nabla\varphi(x_1, \dots, x_n, t) \in C^1([0, T]; H^p)$.

Literature

- [1] Davey A., Stewartson K., On the three-dimensional packets of surfase waves, Proc.R.Soc.London, 1974, Vol.338, Ser.A, pp.101–110.
- [2] Ghidaglia J.-M., Saut J.-C., On the initial value problem for the Davey-Stewartson systems Nonlinearity, 1990, Vol.3, pp.475–506.
- [3] Vakulenko S.A., Justification of asymptotic formula for the perturbed Klein-Fock-Gordon equation, Zap.nauch.semin. LOMI, 1981, Vol.104, pp.84–92.
- [4] Kalyakin L.A., Asymptotic collapse of one-dimensional wave packet in non-linear dispersive medium, Mat.sbornik, 1987, Vol.132, No.4, pp.470–495.
- [5] Ovsjannikov L.V. A nonlinear Cauchy problem in a scale of Banach spaces, Dokl.Akad.Nauk SSSR, 1971, Vol.200, No.4, pp.789–792.